

# Animal Health Care Center

Dr. John B. Dunlap & Dr. Terri Sears  
(540) 943-2273

## Surgical Admission

Date \_\_\_\_\_  
 Patient \_\_\_\_\_ Client \_\_\_\_\_ Wt \_\_\_\_\_ lbs. DR \_\_\_\_\_ LVT \_\_\_\_\_  
 Surgery \_\_\_\_\_ TPR \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_ Contact Number \_\_\_\_\_

	NORM--ABNORM	ABNORMALITIES
General Appearance	N----A	--- Other
Teeth-----	N----A	--- Extras, malocclusion, tartar, g'vitis
Ears-----	N----A	--- Mites, wax, suspicious
Eyes-----	N----A	--- Bloodshot, watery, icteric
Skin-----	N----A	--- Fleas, rash, ticks, flakes, tumors
Cardiovascular-----	N----A	--- Murmurs, arrhythmias, pale mm
Toes-----	N----A	--- Extras- OK to remove? Y__ N__
Hernias-----	N----A	--- Umbilical or Inguinal
Reproductive M/F---	N----A	--- Cryptorchid, had heat, had litter
NPO	NO YES	When? Age?
Seizures-----	NO YES	
<b>TESTING</b>		<input type="checkbox"/> Call for Permission for anything not discussed.
Pap-----	NO YES	<input type="checkbox"/> Permission to do procedures necessary or important.
Ekg-----	NO YES	I understand that even though I may have opted to do pre-anesthetic testing, and AHCC will take all necessary precautions, death under anesthesia does occur and despite our best efforts we make no guarantee of outcome.
Fecal-----	NO YES	
3DX-----	NO YES	
FELV/FIV-----	NO YES	
Tattoo/Microchip _____	NO YES	
Pain Med (TGH)	NO YES	
Flouride Tx	NO YES	
Grooming	NO YES	
Biopsy	NO YES	Owner/Agent Signature: _____

## Surgical release/Post operative care

1. Restrict activity for one week
2. Keep incision clean and dry, no bathing for two weeks
3. Call if any signs of infection occur (swelling, drainage, or pus)
4. If your pet chews and damages the surgical site, please call our office
5. Suture removal: None\_\_ Removal in 10-14 days\_\_
6. Restrict food and water. You may give your pet 1/4 of its regular meal 3-4 times in the next 24 hours. Only give 2-4 oz of water an hour until your pet is satisfied. If vomiting occurs remove food and water for 12 hours and call our office.
7. Coughing can be expected if we intubated. If you notice increase swallowing or coughing you may treat with honey.
8. \_\_\_\_\_

## Declaw

1. Use special pelleted litter provided by AHCC for a complete two-week period, which is very important to reduce the chance of infection
2. Call if any signs of infection, swelling, or bleeding occur
3. Declawed cats must be kept indoors during the healing process
4. Older and/or larger cats may take longer to heal